International Journal of All Research Education and Scientific Methods (IJARESM),
ISSN: 2455-6211, Volume 13, Issue 5, May-2025, Available online at: www.ijaresm.com

Formualtion and Evaluation Processing of Friuts with Special Reference o Actinidia Deliciosa

Mr. Prashant Balasaheb Darade[1], Ms. Maya Jagan Dane[2], Ms. Dube S. B[3], Ms. Dusane G. V.[4], Mr. Rahul S. Mohan[5]

[1,2,3,4,5]Nandkumar Shinde College of Pharmacy, Aghur, Tq. Vaijapur - 423701 ,
Dist Sambhajinagar , Maharshtra

----------**********----------

ABSTRACT

Now a days Jellies are achieving high popularities among the community. The qualitative analysis of market available low quality Jellies are not full filling the nutritive value and appear to be very hazardous to human health. For this very purpose this article is generalized about the full description and nutritive values of jellies along with the methods of making it in homes and all the basic information is gathered from trusted sources online. This article summarize briefly the making of Jellies in home with high nutritive values.

Keywords : Kiwi Frui, Fruit Jelly, Natural Dessert ,Gelatin, Pectin, Healthy Snack, Low-Calorie Dessert, Refreshing Treat, Sugar-Free Jelly

INTRODUCTION

The most vital and healthful component of a balanced diet is fruit. Fruits are a wonderful source of vitamins, enzymes, and minerals, but they are also low in calories, sodium, fat, and cholesterol. Among all fruits, kiwi fruits are the most significant. Kiwi fruits have a wide range of therapeutic uses. It is considered one of the most significant fruits for enhancing immunity due to its antioxidant properties, which aid in the elimination of free radical molecules from the human body. Kiwi fruit's numerous health benefits have caused it to become incredibly popular during the past 20 years.

Actinidia is the genus that produces kiwifruit, which is well-known worldwide for both its tasty flavor and health benefits. In addition to having a distinctive flavor and appearance, the fruit is high in vitamin C. New Zealand is where Kiwi received its name. Kiwi fruit weighs four ounces and is around three inches long. Its flesh is either light green or golden, with rows of tiny, edible black seeds, and its skin is thin, fuzzy, fibrous, sour, but edible. The fruit has a sweet, distinct flavor and a delicate texture. Kiwi fruit can also be eaten in a variety of ways, including fresh, frozen, dried, marmalade, juice, jam, jelly, and nectar. In some cultures, its juice has long been utilized to tenderize meat. Although kiwi season necessitates winter growing, the fruit is excellent for post-harvest processing and storage in both the northern and southern hemispheres. This implies that kiwifruit is available all year round, which is crucial for people who want to regularly eat it for its health advantages. The kiwi berry can be eaten without peeling since it has a soft, edible skin.

Kiwi fruits are rich in phenolics and vitamin C, which contribute to their strong antioxidant activity. After plums, kiwi fruit is the fruit with the second highest antioxidant content that is regularly consumed. [8] Kiwi fruits are rich in phenolics and vitamin C, which contribute to their strong antioxidant activity. After plums, kiwi fruit is the fruit with the second-highest antioxidant content that is regularly consumed. The kiwi fruit is a very nutritious, low-calorie fruit that has the potential to provide significant health advantages. Studies have showed that eating kiwi fruit every day can help prevent and treat conditions including diabetes, asthma, and cough. Compared to citrus fruits or bananas, it contains higher potassium. Numerous investigations have revealed that it includes a wide range of chemicals with potential medical use, including the potential to cure sleep disorders. The kiwi fruit is particularly well-known for its therapeutic uses in China, where it is used to treat rheumatism, cure hemorrhoids, prevent kidney or urinary tract stones, delay the onset of premature graying of the hair, and improve digestion. A variety of beneficial anti- cancer bioactive substances, including prooxidant (at greater concentrations) and antioxidant (at lower concentrations), as well as tumor-specific, cytotoxic, and antibacterial action, are present in kiwi fruit. Because soluble dietary fiber has very few calories, it helps with weight loss and the removal of excess body fat. It also maintains concerns with appetite.

Skin health:
Vitamin C contributes to the production of collagen, which helps to hydrate and supple skin as well as delay the premature emergence of wrinkles. Thus, include kiwis in your diet on a regular basis to help maintain youthful, healthy skin.

Bone health:
For those who are more susceptible to osteoporosis, such as the elderly and expectant mothers, kiwifruit is a great fruit choice. The body has low levels of potassium, phosphate, and calcium, just like in these two target groups. Calcium and phosphorus traces found in kiwis support healthy bones. Additionally, vitamin K is essential for the development of bone mass.

Heart health and blood pressure:
Kiwifruits are high in vitamin C, antioxidants, and phytochemicals, all of which are good for heart health. A study indicated that participants who consumed 100g of kiwifruit every day for eight weeks had significantly higher HDL levels. Additionally, Suksomboon et al. (2019) found that eating two or three kiwifruits a day for 28 days decreased the level of Triacylglycerol by 15% when compared to the control group. Once more, one can monitor elevated salt levels due to the high potassium content. intake, hence maintaining blood pressure management.

Digestive health:
A good source of fiber is kiwifruit. Additionally, actinidin, a unique proteolytic enzyme found only in kiwifruit, aids with digestion by dissolving proteins. Consequently, promoting a seamless digestive system.

Fights insomnia:
Due to its high antioxidant and serotonin content—a hormone essential for mood stabilization and sleep induction—kiwifruit eating can help prevent insomnia.

Diabetes:
With a glycemic index of 39, kiwifruit are classified as "low" GI foods (GI < 55). Not only is it GI, but its accessible carbohydrate content is only around 12%. Once more, the fiber could postpone the breakdown and absorption of carbohydrates.

t is clear that individuals who consume less vitamin C have higher bronchitis and wheezing symptoms, which are more severe in asthmatic and bronchitis vulnerable patients. In older and younger people as well as children, kiwis boost the immune system and lessen the severity of cold and flu-like illnesses. The little kiwi fruit is rich in B6, B12, vitamin C, vitamin K, zinc, fiber, folate, and other elements that work together to promote immunity

Good for vision:
Kiwis guard against the main cause of visual loss, muscle degeneration. The main causes of vision loss are thought to be age, macular degeneration, and retinal degradation. A cup of kiwis contains 220 mg of zeaxanthin and lutein, which help to rebuild the macula and prevent many eye illnesses. Zeaxanthin and lutein are components of the retina and the macula.

Nutritional values of kiwi

Kiwi has a high concentration of nutrients. Kiwi fruit has the highest nutrient content per calorie of any other fruit. It also has a significant quantity of dietary fiber, natural sugar, minerals, and vitamins. It also has zero sodium, low cholesterol, and very little fat. The body receives the ideal nutrients from vitamins and minerals. It keeps the body's digestive system in equilibrium. The following is a discussion of the significant nutritional benefits of kiwi fruit:

VITAMIN C
There are typically three green varieties in every 100g of fresh weight It includes 161.3mg of vitamin C, which is over three times as much as the 80– 120mg of vitamin C included in each 100 g of Sun Gold kiwifruit. than in strawberries and oranges

FOLATE
A good source of dietary folate is kiwifruit. Fresh kiwis are a valuable addition to the diet overall during pregnancy, when it can be challenging to achieve folate requirements because folate in vegetables is rapidly degraded by cooking.

POTASSIUM
Compared to iron and zinc, our bodies require a considerably higher amount of potassium each day. The National Academies Institute of Medicine advises healthy adults to ingest at least 4,700 mg of potassium per day to counteract the effects of salt and to support normal blood pressure levels. Consuming bananas and kiwifruit is a great way to increase potassium. Each 100g of green and gold kiwifruit normally contains 301-315mg of potassium. Kiwi fruit has a salt concentration of only 3 mg per 100g, making it a naturally low-sodium food. Kiwifruit is one of the most balanced fruits in terms of sodium to potassium (Na+/K+) ratio, and its consumption boosts the intake of potassium-rich foods, which can lower blood pressure and hypertension.

DIETARY FIBRE

Particularly, the cell walls of kiwifruit include polysaccharides, which are an important structural component and a source of dietary fiber. It has 2-3% non- starch polysaccharides, which are what give fruits their cell walls and give the diet both soluble and insoluble fibers. The soluble fiber found inside plant cells dissolves in water to generate a gel that slows down the process by which the stomach's contents are expelled. Instead, insoluble fibers, which are derived from the walls of plant cells, give feces weight and keep it moving through the digestive system. 1 gram of insoluble fiber and 0.7 gram of soluble fiber can be found in one large kiwi fruit.

CARBOHYDRATES

One average-sized green kiwi contains 10 grams of carbohydrates, of which 6.2 grams are from naturally occurring sugars and 2.1 grams are from fibers. Kiwis are a fruit with a low glycemic index, with a value of 52. where's requirements may vary, depending on person's gender and age.

Material :

Table No.2

SR.NO.	INGREDIENTS
1	Actinidia deliciosa
2	Stevia
3	Corn starch
4	Gelatin

Actinidia Deliciosa

Fig 1: Actinidia Deliciosa

Scientific name: Actinidia deliciosa Domain: Eukarya

Kingdom: Plantae

Phylum: Magnoliophyte Class: Magnoliopsida Order: Ericales

Family: Actinidiaceae Genus: Actinidia

Species: deliciosa

About: This fruit is considered a miracle since it has an abundance of vitamins, minerals, and phytonutrients. It also has a small amount of sucrose, fructose, and glucose. Due to its high vitamin C and E content, it has the effect of boosting immune function. Numerous pro- and antioxidant-active bioactive compounds that have been shown to have anticancer properties can be found in kiwifruit.

International Journal of All Research Education and Scientific Methods (IJARESM),
ISSN: 2455-6211, Volume 13, Issue 5, May-2025, Available online at: www.ijaresm.com

STEVIA

FIG 2: STEVIA

Sucrose is usually used in homemade jellied fruit products. From a practical point of view, it may be advantageous to substitute other sugars for sucrose, either because of cost, or to reduce the likelihood ofcrystallization, or for flavor modification19. Partial replacement of sucrose with other sugars such as maltose, glucose, syrups, or high fructose corn syrup altered the setting times and certain rheologica properties of model gels20.

CORN STARCH



FIG 3: CORN STARCH

Ingredient, often used to thicken sauces or soups, and to make corn syrup and other sugars. Corn starch is versatile, easily.

2.1 Method :
2.1.1 Formuation Table :

TABLE NO: 3

RMULATION CODE	ACTINIDIA DELICIOSA (gm)	STEVIA (gm)	CORN STARCH (gm)	RESULT
F1	30	5	7	Fail
F2	30	10	4	Success
F3	30	6	7	Fail
F4	30	9	3	Success
F5	30	7	5	Fail
F6	30	9	5	Success
F7	30	10	5	Success

2.1.2 Process :

Collection of fruit sample

↓

Washing

↓

Cutting

↓

Boil in a pot and add corn starch

Addition of Stevia

↓

Packing (filled the hot jelly into clean and dry sterilized jar)

↓

Cooling

↓

Stored in refrigerator

3.Evaluation Test :

FIG 4: KIWI FRUIT JELLY

3.1 Organoleptic Evaluation :

1 Colour : Slightly Yellow wish

2 Odour : Citrusy

3 Taste : Sweet

4 Texture : Smooth

5 Apperance : Good

6 Flavour :

Result and Discussion :

Table No: 4

SR.NO.	PARAMETER	OBSERVATION
1	Color	Slightly yellowish
2	Odour	Citrusy
3	Taste	Sweet
4	Texture	Smooth
5	Consistency	Good

CONCLUSION

A successful kiwi fruit jelly project can demonstrate that kiwi fruit can be a viable and nutritious alternative to traditional jellies, especially when combined with other ingredients like cucumber. The project can highlight the unique sensory appeal and nutritional benefits of kiwi-based jellies, such as higher levels of vitamin C, ascorbic acid, and antioxidants. Additionally, the project can showcase how 3D food printing technology can be used to personalize functional kiwi fruit jellies.

Future Prospective :
The future scope of processing Actinidia deliciosa fruits for jelly preparation is quite promising. As consumer demand for natural and homemade products continues to rise, there is a growing interest in artisanal and gourmet food items, including fruit jellies. In terms of innovation, there are several exciting possibilities. This could result in a wide range of delicious and exotic jelly flavors that cater to diverse palates. Additionally, there is potential for incorporating healthier and natural alternatives to traditional sugar and pectin in jelly recipes. Furthermore, with advancements in food technology, there may be opportunities to develop innovative packaging and preservation methods to extend the shelf life of fruit jellies without compromising their taste and quality. This could involve exploring techniques such as vacuum sealing or modified atmosphere packaging to maintain freshness for a longer duration. Overall, the future of processing Actinidia deliciosa fruits for jelly preparation holds immense potential for creativity, flavor experimentation, and meeting the evolving preferences of consumers. It's an exciting time for fruit jelly enthusiasts and food lovers alike!

REFERENCES

[1]. Akintunde BO., Tunde Akintunde TY., Adejumo AO. (2004) Development of a Manual fruit juice Extracto, NIFOL, 22, 178-182.
[2]. Iwe MO., Obaje PO., Akpapunam MA. (2003) Organoleptic Assessment of Selected Fruit Juices Produced with citrus populace gum powder (Okoho) extracted with the aid of edible Starches. NIFOI, 21, 79-84.
[3]. Fügel R., Carle R., Schieber A. (2005) Quality and authenticity control of fruit purées, fruit preparations and jams – a review. Trends Food Sci. Tech., 16, 433–441.
[4]. Baker RA., Berry N., Hui YH. (1996) Fruit Preserves and Jams. In: Biology, Principles, and Applications Technomic Publishing Co. Inc, Lancaser, PA.
[5]. Egbekun MK., Nda-Suleiman EO., Akinyeye O. (1998) Utilization of fluted pumpkin fruit (Telfairia occidentalis) in marmalade manufacturing. Plant Foods Hum Nutr., 52, 171–176.
[6]. FAO. 1992. Codex Alimentarius, Fruit Juices and Related Products.

[7]. Sindumathi G., Amutha S. (2014) Processing and quality evaluation of coconut based jam. IOSR J. Environ. Sci. Toxicol. Food Technol., 8 (1), 10-14.
[8]. Madhav A., Pushpalatha PB. (2002) Quality upgradation of jellies prepared using pectin extracted from fruit wastes. J. Trop. Agri., 40, 31-34.
[9]. Pramanick P., Zaman S., Bera D., Raha AK., Mitra A. (2014) Mangrove fruit Products: A search for alternative livelihood for Island dwellers of Gangetic delta. Int. J. Pharm. Res. Scholars, 3 (1), 131-137.
[10]. Vidhya R., Narain A. (2011) Formulation and Evaluation of Preserved Products Utilizing under Exploited Fruit, Wood Apple (Limonia acidissima). Am-Eurasian J. Agric. Environ. Sci., 10 (1), 112-118.
[11]. Kapoor M., Beg QK., Bhushan B., Dadhich KS., Hoondal GS. (2000) Production and partial purification and characterization of a thermoalkalistable polygalacturonase from Bacillus sp. MG-cp-2.‖. Process Biochem., 36, 467–473.
[12]. Kashyap DR., Vohra PK., Chopra S., Tewari R. (2001) Applications of pectinases in the commercial sector: A review. Bioresource Technol., 77, 215-227.
[13]. Singh SA., Plattner H., Diekmann H. (1999) Exopolygalacturonate lyase from a Thermophilic
[14]. Bacillus sp. Enzyme Microb. Tech., 25, 420–425.
[15]. Naidu GSN., Panda T. (1998) Production of pectolytic enzymes – a review. Bioprocess Eng., 19, 355-361.
[16]. Hoondal GS., Tiwari RP., Tewari R., Dahiya N., Beg QK. (2002) Microbial alkaline pectinases and their industrial applications: a review. Appl. Microbiol. Biotecnol., 59, 409-418.
[17]. Sriamornsak P. (2003) Chemistry of Pectin and its Pharmaceutical Uses: A Review. Silpakorn University International Journal, 3, 206.
[18]. Ahmed GE. (1981) High methoxyl pectins and their uses in jam manufacture - a literature survey. The British Manufacturing Industries Research Association. Scientific and Technical Surveys No. 127, 16.
[19]. May CD., Stainsby G. (1986) Factors Affecting Pectin Gelation in Wedlock. Williams, P.A. (eds.) Elseveir Applied Science Publishers. New York. 515–523.
[20]. Zhao Y. (2007) In: Berry fruit: Value-Added Products for Health Promotion. CRC press, 448.
[21]. Akintunde BO., Tunde Akintunde TY., Adejumo AO. (2004) Development of a Manual fruit juice Extracto, NIFOL, 22, 178-182.
[22]. Iwe MO., Obaje PO., Akpapunam MA. (2003) Organoleptic Assessment of Selected Fruit Juices Produced with citrus populace gum powder (Okoho) extracted with the aid of edible Starches. NIFOI, 21, 79-84.
[23]. Fügel R., Carle R., Schieber A. (2005) Quality and authenticity control of fruit purées, fruit preparations and jams – a review. Trends Food Sci. Tech., 16, 433–441.
[24]. Baker RA., Berry N., Hui YH. (1996) Fruit Preserves and Jams. In: Biology, Principles, and Applications Technomic Publishing Co. Inc, Lancaser, PA.
[25]. Egbekun MK., Nda-Suleiman EO., Akinyeye O. (1998) Utilization of fluted pumpkin fruit (Telfairia occidentalis) in marmalade manufacturing. Plant Foods Hum Nutr., 52, 171–176.
[26]. FAO. 1992. Codex Alimentarius, Fruit Juices and Related Products. 6.
[27]. Sindumathi G., Amutha S. (2014) Processing and quality evaluation of coconut based jam. IOSR J. Environ. Sci. Toxicol. Food Technol., 8 (1), 10-14.
[28]. Madhav A., Pushpalatha PB. (2002) Quality upgradation of jellies prepared using pectin extracted from fruit wastes. J. Trop. Agri., 40, 31-34.
[29]. Pramanick P., Zaman S., Bera D., Raha AK., Mitra A. (2014) Mangrove fruit Products: A search for alternative livelihood for Island dwellers of Gangetic delta. Int. J. Pharm. Res. Scholars, 3 (1), 131-137.
[30]. http://slism.com/calorie/200339/
[31]. http://www.nutrition-and-you.com/food-nutrition.html
[32]. Vidhya R., Narain A. (2011) Formulation and Evaluation of Preserved Products Utilizing under Exploited Fruit, Wood Apple (Limonia acidissima). Am-Eurasian J. Agric. Environ. Sci., 10 (1), 112-118.
[33]. Kapoor M., Beg QK., Bhushan B., Dadhich KS., Hoondal GS. (2000) Production and partial purification and characterization of a thermoalkalistable polygalacturonase from Bacillus sp. MG-cp-2.‖. Process Biochem., 36, 467–473.
[34]. Kashyap DR., Vohra PK., Chopra S., Tewari R. (2001) Applications of pectinases in the commercial sector: A review. Bioresource Technol., 77, 215-227.
[35]. Singh SA., Plattner H., Diekmann H. (1999) Exopolygalacturonate lyase from a Thermophilic Bacillus sp.

Enzyme Microb. Tech., 25, 420–425.

[36]. Naidu GSN., Panda T. (1998) Production of pectolytic enzymes – a review. Bioprocess Eng., 19, 355-361.

[37]. Hoondal GS., Tiwari RP., Tewari R., Dahiya N., Beg QK. (2002) Microbial alkaline pectinases and their industrial applications: a review. Appl. Microbiol. Biotecnol., 59, 409-418.

[38]. Sriamornsak P. (2003) Chemistry of Pectin and its Pharmaceutical Uses: A Review. Silpakorn University International Journal, 3, 206.

[39]. Ahmed GE. (1981) High methoxyl pectins and their uses in jam manufacture - a literature survey. The British Manufacturing Industries Research Association. Scientific and Technical Surveys No. 127, 16.

[40]. May CD., Stainsby G. (1986) Factors Affecting Pectin Gelation in Wedlock. Williams, P.A. (eds.) Elseveir Applied Science Publishers. New York. 515–523.